

Evaluation of inflammatory markers and HALP score in childhood intussusceptions

Evaluation of inflammatory markers in childhood intussusception

Mustafa Tuşat¹, Sebahattin Memiş²

¹ Department of Pediatric Surgery

² Department of Pediatrics, Faculty of Medicine, Aksaray University, Aksaray, Turkey

Abstract

Aim: Intussusception is the most common abdominal emergency in infancy. This study aims to determine whether HALP score and inflammatory markers scores, such as NLR and PLR, differ between cases requiring surgical reduction and those not requiring surgical reduction in patients diagnosed with intussusception who presented to the pediatric emergency department.

Material and Methods: Our study included 78 children diagnosed with intussusception who presented to the Aksaray Training and Research Hospital Pediatric Emergency Department between January 1, 2018, and September 31, 2024. The cases were divided into two groups based on their treatment method: surgical and non-surgical.

Results: When comparing the groups, a significant difference was found for all three markers ($p < 0.0001$) in terms of neutrophil-lymphocyte ratio (NLR), platelet-lymphocyte ratio (PLR), and HALP score. However, no statistically significant difference was detected for C-reactive protein levels ($p = 0.095$).

Discussion: NLR, PLR, and HALP scores can be utilized as useful markers in evaluating the prognosis of childhood intussusception. The effectiveness of these biomarkers should be validated by future studies.

Keywords

Intussusception, Neutrophil/lymphocyte Ratio, Platelet/lymphocyte Ratio, HALP Score

DOI: 10.4328/ACAM.22763 Received: 2025-05-31 Accepted: 2025-06-30 Published Online: 2025-06-30 Printed: 2025-07-01 Ann Clin Anal Med 2025;16(7):530-534

Corresponding Author: Mustafa Tuşat, Department of Pediatric Surgery, Faculty of Medicine, Aksaray University, Aksaray, Turkey.

E-mail: mustafatusat42@hotmail.com P: +90 382 520 10 20

Corresponding Author ORCID ID: <https://orcid.org/0000-0003-2327-4250>

Other Authors ORCID ID: Sebahattin Memiş, <https://orcid.org/0000-0002-3829-9218>

This study was approved by the Ethics Committee of Aksaray University Health Sciences Scientific Research (Date: 2024-09-05, No:2024/082)

Introduction

Intussusception, a significant cause of acute abdominal emergencies in children, is defined as the telescoping of a proximal bowel segment into a distal one [1]. If left untreated, it disrupts the circulation to the bowel wall, leading to intestinal obstruction, venous congestion, and edema. This can further progress to necrosis, perforation, and shock, [2]. While intussusception primarily affects children under three years of age, it is most commonly observed between five and nine months of age, [3].

While most cases of intussusception in children are idiopathic, approximately 5% of cases have a pathological lead point initiating the intussusception, such as a Meckel's diverticulum, duplication cyst, intestinal polyps, cystic fibrosis, lymphoma, or lymphadenopathy [4].

While asymptomatic transient ileoileal or jejunal intussusceptions often resolve with conservative observation, symptomatic ileocolic intussusceptions in hemodynamically stable patients are primarily managed with pneumatic or hydrostatic reduction. Surgical manual reduction is reserved for cases where intussusception persists after these interventions or when there are signs of generalized peritonitis, perforation, or shock [5].

In recent years, easily measurable markers like the neutrophil-to-lymphocyte ratio (NLR) and platelet-to-lymphocyte ratio (PLR) have been shown to indicate the severity of systemic inflammation and correlate with prognosis in various infectious and inflammatory conditions [6-9]. Recent studies have indicated that the HALP score can serve as a prognostic factor in malignancies affected by inflammation and nutrition. This is attributed to anemia occurring in inflammatory processes, weakened immunity, and severe infections, alongside decreased albumin (a negative acute phase reactant), reduced lymphocyte count, and elevated platelet count [9-11].

In our study, we aim to determine if there is a difference in NLR, PLR, and HALP scores between cases of childhood intussusception requiring surgical manual reduction and those not undergoing surgical treatment. By doing so, we intend to evaluate the effectiveness of these inflammatory markers in predicting the need for surgical manual reduction.

Material and Methods

Study design

Our study was conducted as a retrospective descriptive cohort study. We included 78 children under 18 years of age who presented to the Aksaray University Training and Research Hospital pediatric emergency department with a diagnosis of intussusception and were subsequently followed and treated between January 1, 2018, and September 31, 2024. Children who had previously presented with intussusception, those with an unclear radiological type of intussusception, and those with chronic diseases were excluded from the study.

Data collection

Patient data were obtained from files. The included patients were divided into two distinct groups: a non-surgical group, comprising those treated with conservative observation or hydrostatic reduction, and a surgical group, consisting of patients who did not respond to hydrostatic reduction or

presented with signs of generalized peritonitis, perforation, or shock, subsequently undergoing surgical manual reduction.

For each case, sex, age, prominent complaint at admission (abdominal pain, vomiting, bloody stools, diarrhea), season of admission, ultrasonographically measured length and type of the intussuscepted segment, and C-reactive protein (CRP) level at admission were recorded.

NLR (Neutrophil/Lymphocyte Ratio), PLR (Platelet/Lymphocyte Ratio), and the HALP score $[(\text{Hemoglobin (g/L)} \times \text{Albumin (g/L)} \times \text{Lymphocyte count (n/L)} / \text{Platelet count (n/L)})]$ were evaluated at admission. Additionally, in the surgical group, these inflammatory markers were re-evaluated on postoperative day 3, [12].

Statistical analysis

IBM SPSS (Statistical Package for the Social Sciences) 23 for Windows was used for the statistical analysis of the research data. The Kolmogorov-Smirnov test and skewness and kurtosis values, which are other assumptions of normal distribution, were used for normality tests of numerical variables. Comparisons between two independent groups were analyzed using the Mann-Whitney U test, and comparisons between two dependent groups were analyzed using the Wilcoxon signed-rank test. For descriptive statistics, the median (25-75 Interquartile Range) was used for non-normally distributed data, and numbers and percentages were used for categorical variables. Chi-square and Fisher's Exact tests were used to analyze categorical data. Significance was evaluated based on $p < 0.05$ for each variable.

Ethical approval

This study was approved by the Ethics Committee of Aksaray University Health Sciences Scientific Research (Date: 2024-09-05, No:2024/082).

Results

The study included 78 patients, of whom 37.2% (n=29) were female and 62.8% (n=49) were male, and no significant difference was observed between the groups in terms of gender ($p=0.684$). The median age (25-75IQR) of the patients at the time of admission was 19.0 (13.0-39.75) months, which was 35.0 (13.75-58.50) in the surgical group and 16.5 (12.25-31.0) in the non-surgical group, and age was found to be significantly higher in the surgical group ($p=0.01$) (Table 1). While 71.7% of cases with age ≤ 3 were treated non-surgically, 60.2% of cases with age >3 underwent surgical reduction, and this difference was found to be statistically significant ($p=0.012$).

It was observed that 33.3% (n=26) of the cases presented to the hospital in winter, 30.8% (n=24) in autumn, 24.4% (n=19) in spring, and 11.5% (n=9) in summer. No significant difference was found when comparing the groups based on season ($p=0.349$).

When evaluating the prominent complaints of patients upon admission to the emergency department, it was found that 25% (n=6) of cases presenting with abdominal pain, 90% (n=9) of cases presenting with bloody stools, 47.8% (n=11) of cases presenting with vomiting, and 21.1% (n=4) of cases presenting with diarrhea underwent surgical reduction. A statistically significant difference was observed between the groups in this comparison ($p=0.001$) (Table 1).

When evaluating the groups based on the ultrasound-

detected intussusception type, colocolic and jejunojejunal intussusceptions were not observed in the surgical group. However, 10.3% (n=3) of ileoileal intussusceptions and 60% (n=27) of ileocolic intussusceptions required surgical intervention, with a significant difference found when comparing the groups (p<0.001) (Table 1). Regarding the ultrasonographically measured intussuscepted segment length,

Table 1. Comparison of demographic, clinical characteristics and inflammatory markers of the two groups

	Surgery (n=30)	Non-Surgery (n=48)	P
	MD (25-75 IQR)	MD (25-75 IQR)	
Age (mont)	35.0 (13.75-58.50)	16.5 (12.25-31.0)	0.01*
Invaginated segment (mm)	48 (40-61.25)	20 (15-30)	<0.001*
CRP	9.88 (2.41-21.39)	3.55 (0.60-14.55)	0.095*
NLR	5.50 (4.33-7)	2.01 (1.35-2.52)	<0.001*
HALP score	2.44 (1.85-3.23)	5.58 (4.28-7.37)	<0.001*
PLR	203.37 (168.75-250.79)	101.72 (72.95-131.69)	<0.001*
	Surgery	Non-Surgery	
	n (%)	n (%)	
Gender			
Female	12 (41.4)	17 (58.6)	0.684**
Male	18 (36.7)	31 (63.3)	
Type			
Ileoileal	3 (10.3)	26 (89.7)	<0.001***
Ileocolic	27 (60.0)	18 (40.0)	
Colocolic	0 (0.0)	2 (100.0)	
Jejunojejunal	0 (0.0)	2 (100.0)	
Season			
Spring	8 (42.1)	11 (57.9)	0.349**
Summer	2 (22.2)	7 (77.8)	
Fall	7 (29.2)	17 (70.8)	
Winter	13 (50)	13 (50)	
Clinical symptoms			
Abdominal pain	6 (25)	18 (75)	0.001***
Bloody stool	9 (90)	1 (1)	
Vomiting	11 (47.8)	12 (52.2)	
Diarrhea	4 (21.1)	15 (78.9)	
Fever	0 (0)	2 (100)	
Age (year)			
≤3	15 (28.3)	38 (71.7)	0.012**
>3	15 (60.0)	10 (40.0)	

*Mann-Whitney U test **Pearson Chi-square ***Fisher exact test MD: Median CRP: C-reactive protein, NLR: neutrophil-lymphocyte ratio, PLR: platelet-lymphocyte ratio, HALP: hemoglobin, albumin, lymphocyte, and platelet

Table 2. Comparison of pre-op and post-op day 3 inflammatory markers in surgical treatment group

(n=30)	MD (25-75 IQR)	z	p
NLR-day 1	5.50 (4.33-7.00)	-4.78	<0.001*
NLR-day 3	1.79 (1.63-1.89)		
PLR-day 1	203.37 (168.75-250.79)	-4.78	<0.001*
PLR-day 3	109.71 (91.53-121.50)		
HALP-score day 1	2.44 (1.85-3.25)	-4.78	<0.001*
HALP-score day 3	4.01 (3.34-5.15)		

* Wilcoxon signed-rank test

the median (25-75 IQR) length in the surgical group was 48 (40-61.25) mm, compared to 20 (15-30) mm in the non-surgical group, a statistically significant difference (p<0.001) (Table 1). The median (25-75 IQR) CRP level was 9.88 (2.41-21.39) in the surgical group and 3.55 (0.60-14.55) in the non-surgical group, with no significant difference observed between the groups (p=0.095). The median (25-75 IQR) NLR was 5.50 (4.33-7) in the surgical group and 2.01 (1.35-2.52) in the non-surgical group. The median (25-75 IQR) PLR was 203.37 (168.75-250.79) in the surgical group and 101.72 (72.95-131.69) in the non-surgical group. The median (25-75 IQR) HALP score was 2.44 (1.85-3.23) in the surgical group and 5.58 (4.28-7.37) in the non-surgical group. A statistically significant difference was found for all three markers (NLR, PLR, and HALP score) when comparing the groups (p<0.001) (Table 1). Furthermore, in the surgical group, the median (25-75 IQR) NLR on postoperative day 3 was 1.79 (1.63-1.89), the PLR was 109.71 (91.53-121.50), and the HALP score was 4.01 (3.34-5.15). Consequently, a statistically significant difference was observed for all three markers when comparing their values at admission and on postoperative day 3 (p<0.001) (Table 2).

Discussion

In our study, we found that children diagnosed with intussusception who underwent surgical manual reduction had significantly lower NLR and PLR values and a significantly higher HALP score at admission (p<0.001). Furthermore, we observed that CRP levels, which are commonly used for monitoring and prognosis in numerous inflammatory, infectious, and tissue damage conditions [13, 14], were ineffective in predicting the need for surgical manual reduction (p=0.095). It is generally reported that hematological parameter combinations such as NLR and PLR, calculated from a complete blood count, can be used as biomarkers to predict prognosis in many infectious and inflammatory conditions. This is due to the increase in neutrophil count in the blood and the rise in platelets, which are effective in regulating various inflammatory states in the presence of a systemic infection or inflammatory response [6-8, 12]. In a study of 115 patients who underwent surgical operations for intussusception, with 47 of these cases involving resection, a comparison between the resection and non-resection groups revealed significantly higher NLR and PLR values in the group that underwent resection. This difference was found to be statistically significant, [15]. A retrospective study evaluating the accuracy of NLR in predicting the differentiation between complicated and non-complicated appendicitis, which assessed 81 adult appendicitis cases, reported that an NLR > 7 served as a specific and predictive marker for complicated appendicitis [6]. Another study that evaluated 61 mild and 26 moderate pancreatitis cases found that PLR and NLR values were higher in moderate pancreatitis cases. Statistical analysis revealed a significant difference between the two groups for both NLR and PLR, indicating that NLR and PLR have a discriminatory effect in these cases, [7]. In another study that included 291 appendicitis and 101 control pediatric cases, analysis between the appendicitis and control groups revealed that the NLR value was significantly higher in the appendicitis group. Furthermore, it was reported that the NLR value was significantly higher in

children with gangrenous appendicitis compared to those with acute appendicitis [16]. In our study, we found that both PLR and NLR values were significantly higher in the surgical group. We attribute this to more intense ischemia in the intussuscepted segment, leading to the release of more inflammatory factors into the bloodstream. Additionally, we observed a significant decrease in PLR and NLR values on postoperative day 3 in the surgical group compared to their preoperative values ($p < 0.001$). Research indicates that hematological markers such as albumin, hemoglobin, lymphocytes, and platelets are associated with inflammation or nutritional status. These markers can effectively predict prognosis in inflammatory, infectious, and malignant conditions, [9-11, 17]. A study including 684 cases diagnosed with acute appendicitis reported that the incidence of gangrene, perforation, periappendicular abscess, and postoperative complications was significantly increased in the group with a low HALP score compared to the group with a high HALP score. [11]. In a study evaluating 213 cases in the emergency department, although no statistical significance was found when comparing groups based on albumin, hemoglobin, lymphocyte, and platelet levels, it was reported that patients who underwent surgery due to ileus had significantly lower HALP scores. [18]. In another study that evaluated a total of 436 appendicitis cases, comprising 126 complicated and 310 non-complicated instances, it was reported that the HALP score was significantly lower in the complicated appendicitis group and that the preoperative HALP score could be used to predict the diagnosis [9]. In our study, the median (25-75 IQR) HALP score was found to be 2.44 (1.85-3.23) in the surgical group, while it was 5.58 (4.28-7.37) in the non-surgical group, indicating a significantly lower score in the surgical group ($p < 0.001$). This finding suggests its effectiveness in predicting the need for treatment. We believe this difference is a result of the metabolic response to severe inflammation caused by the compression of the intussuscepted segment in cases requiring surgery.

Studies conducted on children diagnosed with intussusception have identified a positive correlation between the increasing age of the cases and the need for surgical treatment, [19, 20]. A study including 153 intussusception cases in children under 12 years of age reported that the probability of a pathological lead point is very high in children over 5 years old, and early surgical intervention should be considered in these children, [21]. In our study, we also found that age was significantly higher in cases in the surgical group compared to the non-surgical group ($p = 0.01$). We observed that 71.7% of cases aged ≤ 3 years were treated non-surgically, whereas 60.2% of cases aged > 3 years underwent surgical reduction ($p = 0.012$). We believe this is related to the increased incidence of pathologies initiating intussusception with advancing age.

It has been reported that the length of the intussusception segment and the presence of currant jelly stools are effective factors that increase the likelihood of surgical reduction in children with intussusception [19, 20, 22]. In a study investigating successful factors for non-surgical reduction in 99 children diagnosed with intussusception, the average length of the intussuscepted segment in cases requiring surgical treatment was found to be 7.9 cm. In contrast, it was 3.5 cm

in cases that underwent non-surgical reduction. A significant difference was observed between these two groups, and in univariate analysis, bloody stool was identified as an effective predictor for surgical intervention, [20]. In a study investigating factors associated with surgical treatment in 106 children with ileoileal and ileocolonic intussusception, it was reported that the length of the intussuscepted segment > 3.5 mm ($p = 0.042$) and strawberry jelly stool were effective in predicting surgery in the surgical treatment group, [19]. In our study, we also found that 9 out of 10 patients presenting with bloody stools required surgical treatment. Furthermore, the median (25-75 IQR) length of the intussuscepted segment was 48 (40-61.25) mm in the surgical group compared to 20 (15-30) mm in the non-surgical group, a statistically significant difference ($p < 0.001$). We attribute this to the more intense ischemia resulting from obstruction within the longer intussuscepted segment in the surgical group.

Limitation

The main limitations of our study are the retrospective study design and the limited number of cases.

Conclusion

In the treatment of intussusception, early diagnosis and appropriate treatment are vitally important in reducing mortality and morbidity. Alongside ultrasonography, a thorough evaluation of physical examination and clinical findings continues to play a significant role in guiding treatment. Furthermore, NLR, PLR, and HALP scores may contribute to clinical and ultrasonography findings in predicting intussusception cases that require surgical intervention.

Scientific Responsibility Statement

The authors declare that they are responsible for the article's scientific content including study design, data collection, analysis and interpretation, writing, some of the main line, or all of the preparation and scientific review of the contents and approval of the final version of the article.

Animal and Human Rights Statement

All procedures performed in this study were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

Funding: None

Conflict of Interest

The authors declare that there is no conflict of interest.

References

- Guo H, Lei H, Luo J, Yang J, Bian H, Yang H, Guo Q. Clinical manifestation and treatment of intussusception in children aged 3 months and under: a single centre analysis of 38 cases. *BMC pediatrics*. 2025;26:25(1):233.
- Shavit I, Levy N, Dreznik Y, Soudack M, Cohen DM, Kuint RC. Practice variation in the management of pediatric intussusception: a narrative review. *Eur J Pediatr*. 2024;183(11):4897-904.
- Li Y, Zhou Q, Liu C, Sun C, Sun H, Li X et al. Epidemiology, clinical characteristics, and treatment of children with acute intussusception: a case series. *BMC pediatr*. 2023;23(1):143.
- Qian Y, Guo W. Development and validation of a deep learning algorithm for prediction of pediatric recurrent intussusception in ultrasound images and radiographs. *BMC Med Imaging*. 2025;25(1):67.
- Wang S, Wang Y, Jia L, Wang X. Transient and persistent small-bowel intussusception in children: a decision tree analysis model based on ultrasound and clinical findings. *BMC Gastroenterol*. 2025;25(1):294.
- Ali A, Khan GY, Khan MN, Hassan M, Razaq N. Evaluating the diagnostic accuracy of the neutrophil-to-lymphocyte ratio in acute appendicitis: a distinction between complicated and uncomplicated presentations. *Thermsr*. 2025;3(1):1085-91.
- Akoglu EU, Ozdemir S, Ak R, Ozturk TC. The discriminative power of inflammatory markers in patients with mild-to-moderate acute pancreatitis: mean platelet volume, neutrophil-lymphocyte ratio, lymphocyte-monocyte ratio, and neutrophil-monocyte product. *South Clin Istanbul Eurasia*. 2021;32(2):159-65.
- Yin Q, Yin J, Shen L, Zhou Q, Xu W. The early diagnostic value of neutrophil to

- lymphocyte ratio and platelet to lymphocyte ratio in neonatal late-onset sepsis. *Front Pediatr*.2025;13:1483522.
9. Saridas A, Vural N, Duyan M, Guven HC, Ertas E, Cander B. Comparison of the ability of newly inflammatory markers to predict complicated appendicitis. *Open Med*. 2024;19(1):20241002.
 10. Xu H, Zheng X, Ai J, Yang L. Hemoglobin, albumin, lymphocyte, and platelet (HALP) score and cancer prognosis: A systematic review and meta-analysis of 13,110 patients. *Int Immunopharmacol*.2023;114:109496.
 11. Benli S, Tazeoğlu D. The efficacy of hemoglobin, albumin, lymphocytes, and platelets (HALP) score in signifying acute appendicitis severity and postoperative outcomes. *Updates Surg*. 2023;75(5):1197-202.
 12. Demirtas MS, Memis S, Kocagozoglul SG. Evaluation of hemoglobin, albumin, lymphocyte, and platelet score, neutrophil-to-lymphocyte ratio, and platelet-to-lymphocyte ratio in pediatric patients with rotavirus enteritis. *J Clin Pract Res*. 2025;47(1):28-35.
 13. Ural DA, Karakaya AE, Güler AG, Yalçın HS, Tuşat M. Comparative analysis of the acute appendicitis management in children before and during the coronavirus disease-19 pandemic. *KSU Med J*. 2023;18(1):120-5.
 14. Plebani M. Why C-reactive protein is one of the most requested tests in clinical laboratories?. *CCLM*. 2023;61(9):1540-5.
 15. Chen B, Cao J, Yan C, Zheng C, Chen J, Guo C. A promising new predictive factor for detecting bowel resection in childhood intussusception: the lymphocyte-C-reactive protein ratio. *BMC pediatr*.2021;21(1):577
 16. Duran İ, Avci V, Nazik S, Altun E. Neutrophile lymphocyte ratio and platelets lymphocyte ratio in the diagnosis of childhood appendicitis. *J Turk Cli Biochem*. 2017;15(1):1-7.
 17. Tuşat M, Özmen İ, Demirtaş MS, Ateş C, Öztürk AB, Kankılıç NA, et al. Risk factors for mortality and morbidity in Syrian refugee children with penetrating abdominal firearm injuries: an 1-year experience. *TJTES*.2023;29(9):1051.
 18. Çetinkaya HB, Çay F. Usefulness of hemoglobin, albumin, lymphocyte and platelet (HAKP) score in determining the need for surgery in ileus patients in the emergency department: hemoglobin, albumin, lymphocyte and platelet (HALP) score. *Chron Precis Med Res*. 2023;4(3):414-7.
 19. Peyvasteh M, Askarpour S, Ghanavati M, Javaherizadeh H. Factors associated with surgical treatment in pediatric intussusception. *Wien Med Wochenschr*.2022;172(13):313-6.
 20. Issa K, Ali W, AL-Abbas B. Factors associated with success of sonographically guided hydrostatic reduction of ileocolic intussusception in children. *SN Compr Clin Med*. 2021;3(1):242-6.
 21. Banapour P, Sydorak RM, Shaul D. Surgical approach to intussusception in older children: influence of lead points. *J Pediatr Surg*. 2015;50(4):647-50.
 22. Alsinan T, Altokhais T, Alshayeb F, Hajja A, Boukai MA, Alsalameh S, et al. Risk factors for failure of enema reduction of intussusception in children. *Sci Rep*. 2024;14(1):20786.

How to cite this article:

Mustafa Tuşat, Sebahattin Memiş. Evaluation of inflammatory markers and HALP score in childhood intussusceptions. *Ann Clin Anal Med* 2025;16(7):530-534

This study was approved by the Ethics Committee of Aksaray University Health Sciences Scientific Research (Date: 2024-09-05, No:2024/082)